

AD-A161 978

NEW CONFIDENCE INTERVAL ESTIMATORS USING STANDARDIZED  
TIME SERIES(U) CORNELL UNIV ITHACA NY SCHOOL OF  
OPERATIONS RESEARCH AND INDU D GOLDSMAN ET AL

1/1

UNCLASSIFIED

DEC 84 TR-J-84-15 N00014-81-K-0037

F/G 12/1


NL



END

FILED

QTHC



12

AD-A161 978

SCHOOL OF INDUSTRIAL AND  
SYSTEMS ENGINEERING  
GEORGIA INSTITUTE OF TECHNOLOGY  
ATLANTA, GEORGIA 30332

TECHNICAL REPORT NO. J-84-15  
December, 1984

NEW CONFIDENCE INTERVAL ESTIMATORS  
USING STANDARDIZED TIME SERIES

by

David Goldsman

and

Lee Schruben

DTIC  
ELECTE  
DEC 06 1985  
S D E

DTIC FILE COPY

This research was partially supported by the Office of Naval Research under contract N00014-81-k-0037 at Cornell University. Lee Schruben is on the faculty of the School of Operations Research and Industrial Engineering, Cornell University, Ithaca, NY 14853.

This document has been approved  
for public release and sale; its  
distribution is unlimited.

85 12 2 069

### Abstract

We develop new confidence interval estimators for the underlying mean of a stationary simulation process. These estimators can be viewed as generalizations of Schruben's so-called *standardized time series area* confidence interval estimators. Various properties of the new estimators are given.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



In this note, we present new confidence interval estimators for the underlying mean  $\mu$  of a stationary simulation process. These estimators can be viewed as generalizations of the so-called area estimators given in Schruben (1983) and Goldsman (1984).

Consider the stochastic process  $X_1, \dots, X_m$ . Define

$$\bar{X}_j \equiv \frac{1}{j} \sum_{k=1}^j X_k, \quad j=1, \dots, m,$$

$$\sigma^2 \equiv \lim_{m \rightarrow \infty} m \text{Var}(\bar{X}_m), \quad \text{and}$$

$$T_m(t) \equiv \frac{[mt](\bar{X}_m - \bar{X}_{[mt]})}{\sigma\sqrt{m}}, \quad 0 \leq t \leq 1,$$

where  $[.]$  is the greatest integer function.  $\{T_m(t), 0 \leq t \leq 1\}$  is called the *standardized time series*. Schruben (1983) proves that if  $X_1, \dots, X_m$  is a stationary,  $\phi$ -mixing, finite variance sequence of random variables (satisfying one other technical condition), then as  $m \rightarrow \infty$ ,  $T_m(t)$  converges in distribution to a *standard Brownian bridge* process,  $\{B_t, 0 \leq t \leq 1\}$ . Also, the standardized time series is asymptotically independent of  $m\bar{X}_m$ .

Remark: It is well known that  $B_t \sim \text{Nor}(0, t(1-t))$  and  $\text{Cov}(B_{t_1}, B_{t_2}) = \min(t_1, t_2)(1 - \max(t_1, t_2))$ .

Define  $A \equiv \frac{\sigma}{m} \sum_{k=1}^m c_k T_m(k/m)$ , the  $c_k$ 's being pre-specified.

For large  $m$ ,  $A$  is (is approximately distributed as)  $\sigma \text{Nor}(0, V)$ , where

$$V \equiv \frac{1}{2} \sum_{j=1}^m \sum_{k=1}^m \text{Cov}(c_j B_{j/m}, c_k B_{k/m}).$$

It will be computationally convenient to approximate  $V$  by letting  $j = ms$  and  $k = mt$  (so that  $dj dk = m^2 ds dt$ ). Thus,

$$\begin{aligned} V &= \frac{1}{m^2} \sum_{j=1}^m \sum_{k=1}^m c_j c_k \min(j/m, k/m) [1 - \max(j/m, k/m)] \\ &= \int_0^1 \int_0^1 c_{ms} c_{mt} \min(s, t) [1 - \max(s, t)] ds dt \\ &= 2 \int_0^1 \int_0^t c_{ms} c_{mt} s(1-t) ds dt. \end{aligned} \quad (1)$$

So  $A^2/V = \sigma^2 \chi^2(1)$ , and this is called the *weighted area estimator* for the variance  $\sigma^2$ .

Suppose now that we work with the process  $X_1, \dots, X_n$ , where  $n=bm$ , and that this series satisfies Schruben's conditions. Divide the process into  $b$  contiguous batches, each of size  $m$ ; i.e.,  $X_{(i-1)m+1}, X_{(i-1)m+2}, \dots, X_{im}$  comprise batch  $i$ ,  $i=1, \dots, b$ . Each individual batch can be standardized: For  $i=1, \dots, b$  and  $j=1, \dots, m$ , let

$$\bar{X}_{i,j} \equiv \frac{1}{j} \sum_{k=1}^j X_{(i-1)m+k} \quad \text{(average of the first } j \text{ } X\text{'s from the } i\text{-th batch),}$$

$$\bar{\bar{X}}_n \equiv \frac{1}{n} \sum_{k=1}^n X_k \quad \text{(grand mean),}$$

$$T_{i,m}(t) \equiv \frac{[mt](\bar{X}_{i,m} - \bar{\bar{X}}_n)}{\sigma\sqrt{m}}, \quad 0 \leq t \leq 1, \quad \text{and}$$

$$A_i \equiv \frac{\sigma}{m} \sum_{k=1}^m c_k T_{i,m}(k/m).$$

For large enough  $m$ , each of the standardized time series [the  $T_{i,m}(t)$ 's] is approximately distributed as a Brownian bridge; so  $A_i \sim \sigma \text{Nor}(0, V)$ ,  $i=1, \dots, b$ . Further, for large  $m$ , we can treat the batches as if they were (approximately) independent. This yields:

$$\frac{1}{V} \sum_{i=1}^b A_i^2 = \sigma^2 \chi^2(b) .$$

An immediate consequence of Theorem 21.1 of Billingsley (1968) is that

$$Z_n \equiv \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \approx \text{Nor}(0,1) .$$

Then the asymptotic independence result above gives:

$$\frac{Z_n}{\left[ \frac{\sum A_i^2}{\sigma^2 b V} \right]^{1/2}} \approx \frac{\text{Nor}(0,1)}{\left[ \frac{\chi^2(b)}{b} \right]^{1/2}} \sim t(b) .$$

We finally obtain confidence interval estimators for  $\mu$ :

$$\text{Pr} \left\{ \mu \in \bar{X}_n \pm t_{b,1-\alpha/2} \left[ \frac{1}{nbV} \sum_{i=1}^b A_i^2 \right]^{1/2} \right\} \approx 1-\alpha , \quad (2)$$

where  $t_{b,v}$  is the upper- $v$  quantile of the  $t(b)$  distribution.

We consider various choices for the weights (the  $c_k$ 's). These choices and their resulting  $V$ 's [from the integral approximation (1)] are summarized in Table 1.

#### Remarks:

(i) After choosing the weighting sequence  $\{c_j\}$ , the associated  $V$  from Table 1 is used in (2) to form a confidence interval estimator.

(ii) The calculation of  $V$  from (1) is straightforward (but sometimes tedious).

Example: For choice 3 (from Table 1),

$$V = 2 \int_0^1 \int_0^1 \frac{1}{s(1-s)} \frac{1}{t(1-t)} s(1-t) ds dt$$

$$= 2 \int_0^1 \int_0^t \frac{1}{(1-s)t} ds dt = -2 \int_0^1 \frac{\ln(1-t)}{t} dt = \frac{\pi^2}{3}.$$

(iii) The variance estimator which results from choice 1 (the equal weighting case) is asymptotically the same (as  $m \rightarrow \infty$ ) as the so-called area variance estimator from Schruben (1983).

(iv) For each standardized time series,  $\{T_{i,m}^{(m)}\}$ , choice 2 grants greater weight for 'small' values of  $t$ . Choices 3 through 6 give comparatively little weight to the middle ( $t \approx 1/2$ ) of each standardized time series.

(v) Table 2 summarizes an empirical study involving the order 1 exponential autoregressive model [cf. Lewis, (1980)]. The weighted area estimators are seen to perform well for 'large' batch size.

(vi) Denote the random variable corresponding to the half-length of the weighted area estimator by  $H$ . Following Schmeiser (1982) and Goldsman and Schruben (1984) (G-S), it is easy to derive the following:

$$E[H] = \frac{\sigma}{\sqrt{n}} t_{b,1-\alpha/2}^{(2/b)^{1/2}} \frac{\Gamma((b+1)/2)}{\Gamma(b/2)},$$

$$\text{Var}(H) = \frac{\sigma^2}{n} t_{b,1-\alpha/2}^2 \left\{ 1 - \frac{2}{b} \left[ \frac{\Gamma((b+1)/2)}{\Gamma(b/2)} \right]^2 \right\}, \text{ and}$$

The coverage probability,

$$\Pr\{|\bar{X}_n - \mu_1| < H\} = F(t_{b,1-\alpha/2}) - F(-t_{b,1-\alpha/2}), \text{ where}$$

$\Gamma(\cdot)$  is the gamma function and  $F(\cdot)$  is the c.d.f. of the noncentral

of correlation amongst batches which are encountered when using the area estimator.

#### References

Billingsley, P. (1968). *Convergence of Probability Measures*. John Wiley & Sons, New York.

Goldsman, D. (1984). *On Using Standardized Time Series to Analyze Stochastic Processes*. Ph.D. Thesis, School of OR&IE, Cornell University, Ithaca, NY.

Goldsman, D. and L. Schruben (1984). Asymptotic properties of some confidence interval estimators for simulation output. *Management Science*, 30, 1217-1225.

Lewis, P.A.W. (1980). Simple models for positive-valued and discrete-valued time series with ARMA correlation structures. From *Multivariate Analysis-V*, pp. 151-166, P.R. Krishnaiah (ed.) North Holland, Amsterdam.

Schmeiser, B. (1982). Batch size effects in the analysis of simulation output. *Operations Research*, 30, 556-568.

Schruben, L. (1983). Confidence interval estimation using standardized time series. *Operations Research*, 31, 1090-1108.

Table 1 - Choices of Weights and Resulting V's

weighting choice	$c_j, j=1, \dots, m$	V from (1)
1	1	$1/12$
2	$1 - \frac{j}{m}$	$1/45$
3	$\left[ \frac{j}{m} \left( 1 - \frac{j}{m} \right) \right]^{-1}$	$\frac{\pi^2}{3}$
4	$\left  \frac{1}{2} - \frac{j}{m} \right  + \epsilon \quad (\epsilon \geq 0)$	$\frac{1}{320} + \frac{\epsilon}{32} + \frac{\epsilon}{12}$
5	$\left[ \frac{1}{2} - \frac{j}{m} \right]^2 + \epsilon \quad (\epsilon \geq 0)$	$\frac{1}{4032} + \frac{\epsilon}{120} + \frac{\epsilon^2}{12}$

Table 2

Performance of weighted area confidence interval estimators for the mean of an EAR(1) process with coefficient  $\rho = 0.2$  and exponential (mean=1) noise based on 100 independent runs of 2560 observations each. [Choices of weights are summarized in Table 1].

		weighting choice	1	2	3	4( $\epsilon=0$ )	5( $\epsilon=0$ )
		<u>Confidence interval achieved coverage (90% desired)</u>					
b	m						
1	2560		.88	.92	.90	.89	.90
2	1280		.93	.96	.94	.93	.95
5	512		.95	.94	.96	.95	.96
10	256		.93	.93	.94	.95	.96
20	128		.93	.94	.91	.91	.91


		<u>Average confidence interval half-length ( x 10000)</u>				
b	m					
1	2560		1193	1266	1184	1199
2	1280		592	612	598	613
5	512		480	481	486	492
10	256		440	441	436	444
20	128		426	423	415	427

		<u>Sample standard deviation of half-lengths ( x 10000)</u>				
b	m					
1	2560		900	923	876	932
2	1280		297	300	298	294
5	512		153	149	150	150
10	256		103	104	97.5	95.6
20	128		52.6	50.7	57.9	59.5

Table 3 - Typical Small-Sample Values of  $\rho_1$  and  $\rho_2$   
(m = batch size)

$\alpha$	m	$\rho_1$	$\rho_2$
0.5	15	-0.04268	-0.03686
0.5	20	-0.03231	-0.02767
0.5	25	-0.02600	-0.02215
0.0	--	0	0
-0.5	15	0.19445	0.16427
-0.5	20	0.16964	0.14261
-0.5	25	0.14969	0.12550
-0.9	15	0.42568	0.35214
-0.9	25	0.44111	0.36053
-0.9	50	0.43968	0.35646
-0.9	100	0.41352	0.33456

Figure 1: Correlation of  $A_1, A_2$  versus  
batch size  $m$  for an MA(1) process with  $\alpha = -0.5$



# REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS Unrestricted		
2a. SECURITY CLASSIFICATION AUTHORITY Office of Naval Research			3. DISTRIBUTION/AVAILABILITY OF REPORT Unlimited Distribution		
7b. DECLASSIFICATION/DOWNGRADING SCHEDULE Not applicable					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. J-84-15			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Cornell University		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research		
6c. ADDRESS (City, State and ZIP Code) Ithaca, NY 14853			7b. ADDRESS (City, State and ZIP Code) Arlington, VA 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-81-K-0037		
8c. ADDRESS (City, State and ZIP Code) Arlington, VA 22217			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT NO.		
11. TITLE (Include Security Classification) New confidence interval estimators using standardized time series (unclassified)					
12. PERSONAL AUTHOR(S) David Goldsman and Lee Schruben					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM Mar 31 84 TO Aug 25 85		14. DATE OF REPORT (Yr., Mo., Day) 1984 December	
				15. PAGE COUNT 10	
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB GR.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>We develop new confidence interval estimators for the underlying mean of a stationary simulation process. These estimators can be viewed as generalizations of Schruben's so-called standardized time series area confidence interval estimators. Various properties of the new estimators are given.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Lee W. Schruben			22b. TELEPHONE NUMBER (Include Area Code) (607) 256-4856		22c. OFFICE SYMBOL

**END**

**FILMED**

**1-86**

**DTIC**